

Why the phase shifts for solitons on a vortex filament are so large?*

Jan L. Cieřliński[†]

Uniwersytet w Białymstoku, Instytut Fizyki Teoretycznej
ul. Lipowa 41, 15-424 Białystok, Poland

Abstract

The phase “jumps” for solitons interacting on a vortex filament, observed in experiments, have been unaccounted for since more than twenty years. Using explicit formulas describing the interaction of two solitons on a thin vortex filament in the Localized Induction Approximation we show that an appropriate choice of the parameters of the solitons leads to large phase shifts. This result does not depend on the axial flow along the filament.

PACS Numbers: 05.45Yv, 02.30.Ik

Key words and phrases: vortex filament, localized induction equation, solitons, phase shift

*The research supported partially by the KBN grant 2 P03B 126 22.

[†]e-mail: janek@alpha.uwb.edu.pl

It is well known that the equation describing the motion of a single thin vortex filament in an ideal fluid in the localized induction approximation [1] is integrable in the sense of the soliton theory [2]) and admits soliton solutions (known also as kinks or Hasimoto vortices) [3]. It is interesting that the equation itself has been derived as early as 1906 by da Rios and then rediscovered few times, see [4]). However, the experiments carried out some time later have shown that although some soliton-like patterns can be observed and their qualitative behavior agrees with the theory ([5]) but there are some discrepancies concerning quantitative details [6]. Let us mention the ratio of the group velocity and the phase velocity, the dependence between phase velocity, angular velocity and group velocity, and last but not least: the very large “phase jump” during head-on collisions of two solitons [7, 8].

Then, the approximation has been improved by taking into account the axial flow (the flow inside the vortex core, along the filament) [9]. The kinematics of the vortex filament remained integrable but the equation is now more general (it contains one additional free parameter corresponding to the intensity and direction of the axial flow) [9, 10]:

$$\mathbf{r}_{,t} = \alpha(\mathbf{r}_{,s} \times \mathbf{r}_{,ss}) + \beta(\mathbf{r}_{,sss} + \frac{3}{2}(\mathbf{r}_{,ss})^2 \mathbf{r}_{,s}) , \quad (\mathbf{r}_{,s})^2 = 1 , \quad (1)$$

where $\mathbf{r} = \mathbf{r}(s, t) \in \mathbf{R}^3$, t is the time and s is arc length parameter along the vortex filament, finally α and β are assumed be constant ($\alpha > 0$). The case $\beta = 0$ corresponds to Localized Induction Equation of da Rios, closely associated with the cubic nonlinear Schrödinger equation and continuum Heisenberg ferromagnet model (see [11]). The case $\alpha = 0$ is associated with the modified Korteweg-de Vries equation. Usually α, β are treated as parameters of the theoretical model and they have to follow from experimental data.

The exact formulas for multi-soliton solutions can be obtained using different methods (compare [3, 9, 10, 12]). We follow the soliton surfaces approach [11, 13] as most convenient. The one-soliton solution is obtained without difficulties [13, 14] but two-soliton solution looks much more complicated (compare [12, 15]). An essential simplification of the two-soliton formula is given in [16, 17].

However, the following statement written more than 16 years ago seems to be still accepted: *“The shapes of the ‘Hasimoto soliton’ seem adequate to fit experimental data, but the ratio between group velocity and phase velocity for the wave is a constant (2.0) in LIA, whereas the experiments show a dependence on the ratio of wavelength to core size. Also unexplained is a*

large phase advance on collision of two experimental ‘solitons’. An intriguing attempt to go beyond LIA as originally conceived was presented by Fukumoto & Miyazaki. Their proposal to include an arbitrary axial flow (...) leaves the shapes of the solitons unchanged, changes the velocity of the propagation, but, unfortunately, does nothing for the troublesome phase jump.” [6].

The main goal of the present paper is to show that the large phase jumps are not longer “troublesome” and are in good agreement with the theory.

In order to describe the interaction of two solitons on a single vortex filament we use the following exact formula for two-soliton solution:

$$\mathbf{r}_2 = \frac{1}{2K} \begin{pmatrix} d_1 (e^{Q_2} \cos P_1^+ + e^{-Q_2} \cos P_1^-) + d_2 (e^{Q_1} \cos P_2^+ + e^{-Q_1} \cos P_2^-) \\ d_1 (e^{Q_2} \sin P_1^+ + e^{-Q_2} \sin P_1^-) + d_2 (e^{Q_1} \sin P_2^+ + e^{-Q_1} \sin P_2^-) \\ d_+ \sinh(Q_1 + Q_2) + d_- \sinh(Q_1 - Q_2) + d_0 \sin(P_1 - P_2) - 2sK \end{pmatrix} \quad (2)$$

where

$$K = \cosh Q_1 \cosh Q_2 \cosh \Delta + \sinh Q_1 \sinh Q_2 \sinh \Delta + \sinh \Delta \cos(P_1 - P_2) ,$$

$$P_k^\pm = P_k \pm \delta_k \quad (k = 1, 2) ,$$

$$Q_k = 2s \operatorname{Im} \lambda_k + 2t \operatorname{Im} \omega_k + Q_{k0} ,$$

$$P_k = -2s \operatorname{Re} \lambda_k - 2t \operatorname{Re} \omega_k + P_{k0} ,$$

$$d_\pm = (d_1 \pm d_2) e^{\pm \Delta} , \quad d_0 = d_1 \sin \delta_1 = -d_2 \sin \delta_2 , \quad d_k = \frac{\operatorname{Im} \lambda_k}{|\lambda_k|^2}$$

$$\omega_k := \omega(\lambda_k) , \quad \omega(\lambda) = -2\alpha \lambda^2 - 4\beta \lambda^3 ,$$

(by the way: changing the function $\omega(\lambda)$ we get other equations of a large hierarchy of integrable systems, compare [18]), λ_k are complex parameters, and, finally, Q_{k0} , P_{k0} , δ_1 , δ_2 , Δ are real constants.

The constants Q_{k0} , P_{k0} are arbitrary but not very important, they just fix the initial positions of the solitons. The constants δ_1 , δ_2 , Δ depend on λ_k as follows:

$$e^{\Delta + i\delta_1} := \frac{(\lambda_1 - \lambda_2) \bar{\lambda}_2}{(\lambda_1 - \bar{\lambda}_2) \lambda_2} , \quad e^{\Delta + i\delta_2} := \frac{(\lambda_2 - \lambda_1) \bar{\lambda}_1}{(\lambda_2 - \bar{\lambda}_1) \lambda_1} . \quad (3)$$

Two complex parameters λ_1, λ_2 can be replaced by four real parameters a_1, a_2, b_1, b_2 (where $\lambda_k = a_k + ib_k$), or, which is even more convenient, by c_k, d_k expressed by:

$$c_k = \frac{a_k}{a_k^2 + b_k^2}, \quad d_k = \frac{b_k}{a_k^2 + b_k^2}, \quad a_k = \frac{c_k}{c_k^2 + d_k^2}, \quad b_k = \frac{d_k}{c_k^2 + d_k^2}. \quad (4)$$

One can also use the parameter T_k (compare [19]) responsible for the shape of the soliton: $T_k := c_k/d_k \equiv a_k/b_k$. The parameter Δ depends on c_k, d_k as follows:

$$\Delta = \frac{1}{2} \ln \frac{(d_1 - d_2)^2 + (c_1 - c_2)^2}{(d_1 + d_2)^2 + (c_1 - c_2)^2} \quad (5)$$

The two-soliton formula (2) is obtained in the framework of the soliton surfaces approach [16, 17]. Its form is relatively simple (especially when compared with other, equivalent, formulas, see [9, 10, 14, 15]) and implies a lot of interesting consequences. Formulas for N -soliton solution can be found for instance in [18] (see also [9, 10, 17]).

Let us first discuss the one-soliton solution (in the case $\omega(\lambda) = -2\lambda^2$ found by Hasimoto [3], extended for an arbitrary analytic $\omega(\lambda)$ by Sym [13]):

$$\mathbf{r}_1 = \begin{pmatrix} 0 \\ 0 \\ -s \end{pmatrix} + \frac{d_1}{\cosh Q_1} \begin{pmatrix} \cos P_1 \\ \sin P_1 \\ \sinh Q_1 \end{pmatrix}. \quad (6)$$

Physical characteristics of the single soliton solution $\mathbf{r}_1$ can be computed in the standard way. The amplitude of the Hasimoto vortex (i.e., its maximal distance from the z -axis) is given simply by d_1 . The maximum of the wave envelope ($Q_1 = 0$) performs a helical movement and its z -coordinate (z_{max1}) moves at a constant speed (group velocity v_1^g):

$$v_1^g = \dot{z}_{max1}(t) = \frac{\text{Im}\omega_1}{\text{Im}\lambda_1} = -4(\alpha a_1 + \beta(3a_1^2 - b_1^2)). \quad (7)$$

The solitons can be classified into 3 classes: lump solitons ($|a_1| > |b_1|$), cusp solitons ($|a_1| = |b_1|$) and loop solitons ($|a_1| < |b_1|$) [10]. The names correspond to the shape of the curve $z = z(\rho)$ (where $\rho = \sqrt{x^2 + y^2}$). In experiments only the lump solitons were observed. If the vortex filament has some loops, then the localized induction approximation is not longer valid and these additional interactions probably can destroy the loops. However,

we remark that in the case of smoke rings (where such interactions are much more important) the localized induction equation was successfully applied yielding some patterns which can be observed experimentally ([20], compare [21]).

The size of the lump soliton can be estimated by its half-width. The half-width D_1 is defined as a difference between the two values of z corresponding to Q_1 satisfying $\cosh Q_1 = 2$:

$$D_1 = d_1 \left(-\sqrt{3} + \left(1 + \left(\frac{a_1}{b_1} \right)^2 \right) \ln(2 + \sqrt{3}) \right) \quad (8)$$

Another possibility is to estimate the wavelength of the soliton (i.e., the distance along the z -axis between neighboring wavecrests)

$$\Lambda_1 \approx \frac{\pi}{a_1} . \quad (9)$$

Not entering into technical details we mention that more precise computations lead to the result $\Lambda_1 = \pi/a_1 + d_1(\tanh Q'_1 - \tanh Q''_1)$, where Q'_1 and Q''_1 depend on s and t . The wavelength Λ_1 is not constant along the filament and depends on t . In experiments usually the central wavelength is measured. The wavelength Λ_1 equals π/a_1 with an accuracy $2d_1/\Lambda_1$ which is less than $1/\pi$. Usually, for $d_1 \ll \Lambda_1$, the accuracy is much better and, therefore, Λ_1 is treated as a constant value given by (9). Note that always $\Lambda_1 \geq 2\pi d_1$. Obviously, the formulas (7),(8),(9) are valid for the second soliton ($k = 2$) as well.

The formula (2) enables us to compute the asymptotic behavior of $\mathbf{r}_2$. Assuming $v_1^g \neq v_2^g$ we consider the limit $Q_2 \rightarrow \pm\infty$ (or, more precisely, $|Q_1| \ll |Q_2|$). Thus

$$\mathbf{r}_2 \xrightarrow{Q_2 \rightarrow \pm\infty} = \begin{pmatrix} 0 \\ 0 \\ \pm d_2 - s \end{pmatrix} + \frac{d_1}{\cosh(Q_1 \pm \Delta)} \begin{pmatrix} \cos(P_1 \pm \delta_1) \\ \sin(P_1 \pm \delta_1) \\ \sinh(Q_1 \pm \Delta) \end{pmatrix} \quad (10)$$

The shape and the velocity of the soliton do not change during the interaction. The only result of the interaction is the phase shift. In fact we have two phase shifts: the shift Δ_1^{ph} along $\mathbf{e}_3$ axis and the shift of the angular variable P_1 . The first one is much more important and can be measured in experiments. Analogical considerations are valid in the limit $Q_1 \rightarrow \pm\infty$.

Let us consider the case of the head-on collision of two solitons (this case was observed in experiments [8]). We assume $v_1^g < 0$, $v_2^g > 0$ (head-on collision) and also $b_k > 0$ (a technical assumption which can be made without loss of the generality). We are going to compute the phase shift of the first soliton (indexed by 1) after the interaction with the second soliton. If $t \rightarrow \pm\infty$, then $Q_2 \rightarrow \pm\infty$ and the z -coordinate of the envelope maximum for the first soliton in asymptotic regions ($Q_2 \rightarrow \pm\infty$) can be easily derived from (10):

$$z_{max1}^\pm = v_1^g t + \frac{Q_{10}}{2b_1} \pm d_2 \pm \frac{\Delta}{2b_1} \quad (11)$$

The phase shift $\Delta_1^{ph} := (z_{max1}^+ - z_{max1}^-) \text{sgn}(v_1^g)$ and, taking into account $v_1^g < 0$, we have:

$$\Delta_1^{ph} = -2d_2 - \frac{\Delta}{b_1} = -2d_2 + \frac{c_1^2 + d_1^2}{2d_1} \ln \frac{(d_1 + d_2)^2 + (c_1 - c_2)^2}{(d_1 - d_2)^2 + (c_1 - c_2)^2}. \quad (12)$$

Analogical considerations lead to the similar formula for the phase shift of the second soliton: $\Delta_2^{ph} = -2d_1 - \Delta/b_2$.

To compute the phase velocity of a soliton we have to determine positions of the individual wave peaks. Considering the projection of the soliton on the xz -plane we have to solve (at least implicitly) the equation $\frac{\partial}{\partial s} \left(\frac{\cos P_k}{\cosh Q_k} \right) = 0$. The resulting velocity of a wavecrest depends on its position Q_1 :

$$v_k^{ph} = 2\alpha \frac{b_k^2 - a_k^2}{a_k} + 4\beta(3b_k^2 - a_k^2) - \frac{2b_k^2(\alpha + 4\beta a_k)}{a_k \cosh^2 Q_k} \left(3 - \frac{2b_k^2}{(a_k^2 + b_k^2) \cosh^2 Q_k} \right). \quad (13)$$

In the analysis of solitons on a vortex filament the phase velocity has been always assumed to be constant sufficiently far from the envelope maximum ($Q_k \rightarrow \infty$). However in experiments usually the central region of the kink (i.e., $Q_k \approx 0$) can be studied with the best accuracy. Anyway, the theoretical treatment of the phase velocity is relatively difficult and have to be made with care. Experimental results concerning the phase velocity are also “*rather crude*” [7, 8].

Another quantity estimated from experimental observation is the rate of rotation of the filament maximum (Ω_k), namely $P_k(t)|_{Q_k=0} = \Omega_k t + \text{const}$, where

$$\Omega_k = -4(\alpha + 4\beta a_k)(a_k^2 + b_k^2). \quad (14)$$

To compare the presented theory with experimental results we recall that the two-soliton solution is parameterized by only 4 essential parameters a_1, b_1, a_2, b_2 (the parameters Q_{k0}, P_{k0} , related to initial data, are not important for our analysis) and two more parameters (α, β) characterize the physical model. In experiments one can measure more or less precisely much more quantities: amplitudes (d_k), torsions, velocities (v_k^g, v_k^{ph}), widths (D_k, Λ_k), angular velocities (Ω_k), phase shifts (Δ_k^{ph} and angular phase shifts) etc.

We choose d_k and D_k (or Λ_k) as fundamental quantities. They can be measured with reasonable accuracy. They are very convenient in theoretical analysis because from (8) we can express c_k in terms of d_k and D_k :

$$c_k = \pm d_k \left(\sqrt{\frac{\frac{D_k}{d_k} + \sqrt{3}}{\ln(2 + \sqrt{3})}} - 1 \right). \quad (15)$$

and all other characteristics of the solution are expressed by c_k, d_k (by virtue of (4) we can use the variables c_k, d_k instead of a_k, b_k). From (9) we can also express c_k by Λ_k :

$$\frac{c_k}{d_k} = \frac{\Lambda_k}{2\pi d_k} + \sqrt{\left(\frac{\Lambda_k}{2\pi d_k}\right)^2 - 1}, \quad (16)$$

where $c_k > 0$ is assumed. Combining (15) and (16) we get

$$D_k = \left(2.63\xi_k \left(\xi_k + \sqrt{\xi_k^2 - 1} \right) - 1.73 \right) d_k, \quad (17)$$

where $\xi_k := \Lambda_k/(2\pi d_k)$ (we recall that $\xi_k \geq 1$).

According to the typical experimental situation we assume $d_1 \approx d_2$ and denote $d := d_1$. Then

$$\Delta_k^{ph} \approx (-2 + 0.66\tilde{\Delta})d + 0.38\tilde{\Delta}D_k \quad (18)$$

where $\tilde{\Delta} := \ln(1 + 4d^2(c_1 - c_2)^{-2})$.

From (17) it follows $D_k > \Lambda_k$ for $\Lambda_k/d_k > 10$ and $D_k > 2\Lambda_k$ for $\Lambda_k/d_k > 17$. Moreover, $\tilde{\Delta} > 1$ if $|c_1 - c_2| < 1.53d$, which can be estimated as $|\Lambda_1 - \Lambda_2| < 5d$. Then $\Delta_k^{ph} \sim 0.4D_k$ but this relation can easily “improved” (e.g., $\Delta_k^{ph} \sim D_k$) if solitons are wider and more similar to each other. We arrive at the conclusion that head-on collision of lump solitons of similar size has to result in large phase shifts of both solitons.

The experimental observations of the soliton interactions have shown that *“the forward advance (...) is of the order of the central wavelength of the isolated kink wave”* ([8], compare Fig. 4). The analysis presented above is in perfect agreement with this observation. For $\tilde{\Delta} \approx 3$ we have a very interesting coincidence: the phase shift is equal to the halfwidth of the soliton.

We stress a very important point: the formulas (12),(18) are valid for any α, β . Therefore our conclusion does not depend on the parameters of the physical model (provided that we confine ourselves to the localized induction approximation (1), of course), actually we can take even more general integrable equation described by an arbitrary function $\omega(\lambda)$.

We propose the following procedure of comparing the presented theory with experimental data. First, taking d_k and D_k close to experimental values we use (8) to compute c_1, c_2 (see (15)). In order to get a large phase shift we need $|c_1 - c_2|$ to be rather small, therefore c_1 and c_2 should be of the same sign (this sign is related to the sign of α , which in turn depends on the the direction of the vorticity field along the filament, we take $\alpha > 0$ [8]). In any case we can use (12) to choose right signs of c_k . Then, we use (7) to compute v_1^g, v_2^g . Thus we have a system of two linear algebraic equations for α and β . The system is invariant with respect to the transformation: $\alpha \rightarrow -\alpha, c_k \rightarrow -c_k$ which means that an appropriate signs of c_k yield $\alpha > 0$. Finally, having all six parameters, we may compare other quantities (e.g., rotation rates and phase velocities of the solitons) with experiment. Some quantities (e.g., phase shifts) need only 4 parameters to be computed (see above).

Konno and Ichikawa [10], trying to fit their two-soliton solution to the experimental results of [8], explained very well amplitudes and group velocities of both solitons, to some extent their widths, and also the angular velocity and the phase velocity of one of the solitons (the corresponding experimental data for the second soliton were not reported in [8]). We are able to explain (with reasonable accuracy) the same data but also both phase shifts.

Experimental data from [7, 8] (in units based on centimeter, second and radian) for the first soliton read as follows: $d_1 = 0.49, v_1^g = -33, \Lambda_1 = 6.8, \Omega_1 = -2.4, v_1^{ph} = -30$, the torsion $\tau_1 \equiv 2a_1 = 0.89$. The second soliton is described with lower precision: $d_2 \sim d_1, v_2^g \approx -v_1^g, 2\Lambda_2 \sim \Lambda_1$. Finally, Δ_k^{ph} is “of the same order” as Λ_k . My rough estimate, based on [8] (Fig. 4), is: $v_2^g = 30, \Lambda_2 \sim 3.5, \Delta_1^{ph} \sim 5, \Delta_2^{ph} \sim 1.5$.

Performing the above procedure we get the following parameters: $a_1 = 0.46, b_1 = 0.11, a_2 = 0.78, b_2 = 0.28$. Then we assume $v_1^g = -33, v_2^g = 30$ and finally we get: $d_1 = 0.49, d_2 = 0.41, \Lambda_1 = 6.8, \Lambda_2 = 4.0, v_1^{ph} = -30.5,$

$$\Omega_1 = -2.5, \Delta_1^{ph} = 5.2, \Delta_2^{ph} = 1.4.$$

The agreement with experimental data is quite good (especially for the first soliton). Trying to get d_2 closer to the value 0.49 we noticed an essential decrease of Δ_2^{ph} or increase of Ω_1 . Taking $d_2 = 0.41$ is a kind of compromise.

In this paper I revisited the Localized Induction Equation to show that, in spite of popular views, that model explains interactions of solitons observed in experiments including the existence of the large phase shifts. A little bit surprising is that my result is equally true for the Localized Induction Equation with and without axial flow.

The theoretical investigation of the Localized Induction Equation, carried out in many directions (compare [9, 10, 17, 22, 23, 24]), usually are not easy to be used by experimentalists. I hope that the results of the present paper enable much better analysis of existing experimental results (which seem to be published only fragmentarily) and will prompt further experimental work in this field. Of special interest should be any observations of the interaction of vortex solitons in the superfluid helium because such vortices are really thin which is assumed in the localized induction approximation.

Acknowledgements. The research was partially supported by the Polish Committee for Scientific Research (KBN grant No. 2 P03B 126 22).

References

- [1] G.K.Batchelor: *An Introduction to Fluid Dynamics*, Cambridge Univ. Press 1967.
- [2] G.L.Lamb: *Elements of Soliton Theory*, Wiley & Sons, New York 1980.
- [3] H.Hasimoto: *A soliton on a vortex filament*, *J. Fluid Mech.* **51** (1972), 477-485.
- [4] R.L.Ricca: *Rediscovery of da Rios equation*, *Nature* **352** (1991), 561-562.
- [5] E.J.Hopfinger, F.K.Browand: *Vortex solitary waves in a rotating turbulent flow*, *Nature* **295** (1982), 393-395.
- [6] H.Aref, T.Kambe: *Report on the IUTAM symposium: fundamental aspects of vortex motion*, *J. Fluid Mech.* **190** (1988), 571-595.
- [7] T.Maxworthy, E.J.Hopfinger, L.G.Redekopp: *Wave motions on vortex cores*, *J. Fluid Mech.* **151** (1985), 141-165.
- [8] T.Maxworthy, M.Mory, E.J.Hopfinger: *Waves on vortex cores and their relation to vortex breakdown*, [in:] *Proceedings of AGARD Conference on Aerodynamics of Vortical Type Flows in Three Dimensions* (1983), AGARD CPP-342, paper 29.
- [9] Y.Fukumoto, T.Miyazaki: *Three-dimensional distortions of a vortex filament with axial flow*, *J.Fluid Mech.* **222** (1991), 369-416.
- [10] K.Konno, Y.H.Ichikawa: *Solitons on a Vortex Filament with Axial Flow*, *Chaos, Solitons & Fractals* **2** (1992), 237-250.
- [11] A.Sym: *Soliton Surfaces II. Geometric Unification of Solvable Nonlinearities*, *Lett. Nuovo Cim.* **36** (1983), 307-312.

- [12] K.Konno, M.Mituhashi, Y.H.Ichikawa: *Soliton on Thin Vortex Filament*, *Chaos, Solitons & Fractals* **1** (1991), 55-65.
- [13] A.Sym: *Soliton surfaces and their applications*, [in:] *Geometric Aspects of the Einstein Equations and Integrable Systems* (Lecture Notes in Physics N. 239), pp. 154-231, R.Martini (ed.), Springer-Verlag, Berlin 1985.
- [14] A. Sym: *Soliton surfaces VI. Gauge Invariance and Final Formulation of the Approach*, *Lett. Nuovo Cim.* **41** (1984), 353-360.
- [15] D.Levi, A.Sym, S.Wojciechowski: *N-Solitons on a Vortex Filament* *Phys. Lett. A* **94** (1983), 408-411.
- [16] J.Cieřliński: *Two solitons on thin vortex filament*, *Phys. Lett. A* **171** (1992), 323-326.
- [17] J.Cieřliński: *The Darboux-Bianchi-Bäcklund transformation and soliton surfaces*, [in:] *Nonlinearity and Geometry*, Polish Scientific Publishers PWN, Warsaw 1998.
- [18] J.Cieřliński: *An effective method to compute N-soliton Darboux matrix and N-soliton surfaces*, *J. Math. Phys.* **32** (1991), 2395-2399.
- [19] E.J.Hopfinger, F.K.Browand, Y.Gagne: *Turbulence and waves in rotating tank*, *J. Fluid Mech.* **125** (1982), 505-534.
- [20] J.Cieřliński, P.K.H.Gragert, A.Sym: *Exact Solution to Localized-Induction-Approximation Equation Modelling Smoke Ring Motion*, *Phys. Rev. Lett.* **57** (1986), 1507-1510.
- [21] A.J.Chorin: *Hairpin removal in vortex interactions II*, *J. Comput. Phys.* **107** (1993), 1-9.
- [22] J.Langer, R.Perline: *Poisson Geometry and the Filament Equation*, *J. Nonlinear Sci.* **1** (1991), 71-93.
- [23] K.Suzuki, T.Ono, T.Kambe: *Riemannian Geometrical Analysis of the Motion of a Vortex Filament: A System of $C^\infty(S^1, SO(3))$* , *Phys. Rev. Lett.* **77** (1996), 1679-1682.
- [24] T.Nishiyama: *On the motion of a vortex filament in an external flow according to the localized induction equation*, *Proc. Royal Soc. Edinburgh* **129A** (1999), 617-626.